

On The Problem with Only Zeroth Landau Level Occupancy of Electrons in Strongly Magnetized Neutron Star Matter in β -equilibrium Condition- The Role of Magnetic Dipole Moments

Sutapa Ghosh^{a)}, Soma Mandal^{a)} and Somenath Chakrabarty^{b)}

^{a)}Department of Physics, University of Kalyani, Kalyani 741 235, India

^{b)}Department of Physics, Visva-Bharati, Santiniketan 731 235, India, E-mail:somenath@vbphysics.net.in

It is shown explicitly that if only the zeroth Landau level is occupied by electrons in strongly magnetized neutron star matter in β -equilibrium, then both the proton and neutron matter sectors become spin symmetric. Whereas, the study of Pauli para-magnetism of neutron matter sector shows that such a scenario is physically impossible. It is also shown that in dense stellar hadronic matter in β -equilibrium in presence of strong quantizing magnetic field, with $\sigma - \omega - \rho$ exchange type mean field interaction and with the inclusion of magnetic dipole moments does not allow electrons to occupy only the zeroth Landau level.

PACS numbers: 97.60.Jd, 97.60.-s, 75.25.+z

The study of dense stellar matter in presence of strong magnetic field, which is one of the oldest subject of physics, has gotten a new life after the discovery of a few magnetars, which are believed to be strongly magnetized young neutron stars [1, 2, 3, 4]. The surface magnetic field of these exotic objects are observed to be $\geq 10^{15}$ G. It is then quite possible that the field at the core region can go up to 10^{18} G. Although the origin of such strong magnetic field is not known till today, the studies on the effect of such strong magnetic field on various physical properties and processes in neutron stars particularly at the core region are extremely interesting and important fields of research in nuclear astrophysics as well as in high energy astrophysics [5, 6, 7, 8, 9, 10]. Not only that, the general relativistic studies of such objects have also got some importance [11, 12, 13]. The studies of physical properties of magnetically deformed stellar objects with general relativity are found to be extremely interesting.

An extensive studies have already been done on the equation of state of strongly magnetized neutron stars [5, 6, 9, 10]. The effect of strong magnetic field on the physical processes, e.g., the weak reactions and decays have also been studied in neutron star matter [7, 8]. As a consequence the cooling rate of neutron stars by neutrino emission also change significantly in presence of strong magnetic fields. It is also found that the effect of strong magnetic field on all these physical aspects of neutron stars become most prominent if the charged particles (e.g., electron, proton, etc.) occupy only their zeroth Landau levels.

Now it is known that the maximum value of Landau quantum number occupied by electrons at zero temperature is given by

$$n_{\max}^{(e)} = \frac{[(\mu_e^2 - m_e^2)]}{2eB} \quad (1)$$

where $[]$ indicates the integer part only. Obviously, this maximum value is a function of both magnetic field strength and density of stellar matter through μ_e , the electron chemical potential, m_e and e are the corresponding rest mass and the magnitude of charge. Here we assume the gauge $A^\mu \equiv (0, 0, xB, 0)$, so that the magnetic field B is along x-direction and is constant. Then it is very easy to show, that for electrons, this quantity becomes zero if $B \sim 10^{17}$ G in dense stellar matter of astrophysical interest.

The aim of this article is to show that if the magnetic field in a neutron star is high enough so that the electrons occupy only their zeroth Landau level, then there will be a serious problem on the existence of predicted spin symmetric neutron and proton components in dense neutron star matter. We shall also show that the same conclusion is also true for interacting hadronic matter with non-zero magnetic dipole moments of the constituents, including the anomalous magnetic moment of neutron. In fig.(1) we have shown how $n_{\max}^{(e)} = 0$ value for electrons depends on the strength of magnetic field and density of stellar matter. The curve in this figure indicates $n_{\max}^{(e)} = 1$ for electrons as a function of magnetic field strength and matter density. The space above the line is for $n_{\max}^{(e)} = 0$, whereas below this line, all possible non-zero allowed values. The curve therefore separates the region where only zeroth Landau level is populated from the region where other non-zero Landau levels are occupied by electrons.

To sketch the curve as shown in fig.(1), we set $n_{\max}^{(e)} = 1$ then it is very easy to show that

$$\mu_e = m_e(1 + 2h_f)^{1/2} \quad (2)$$

where $h_f = B/B_c^{(e)}$ and $B_c^{(e)} = 4.41 \times 10^{13}$ G, the quantum mechanical limit of magnetic field strength for electrons (this is the typical strength of magnetic field at which the value of cyclotron quantum for electron exceeds the

corresponding rest mass energy). Hence we can write down for the electron density assuming that they are moving freely within the system

$$n_e = \frac{eB}{2\pi^2} \sum_{n=0}^{n_{\max}^{(e)} (=0)} (2 - \delta_{n0})(\mu_e^2 - m_e^2 - 2neB)^{1/2} = n_p \quad (\text{from charge neutrality condition}) \quad (3)$$

Again the proton density is given by (here also for the sake of simplicity we have not considered interaction)

$$n_p = \frac{eB}{2\pi^2} \sum_{n=0}^{n_{\max}^{(p)}} (2 - \delta_{n0})(\mu_p^2 - m_p^2 - 2neB)^{1/2} \quad (4)$$

where μ_p and m_p are respectively the chemical potential and the mass of proton. Now substituting eqn.(4) into eqn.(3), we can solve numerically for μ_p for a given B . Then assuming β -equilibrium condition, we have $\mu_n = \mu_p + \mu_e$, the neutron chemical potential. We assume that neutrinos / anti-neutrinos are non-degenerate, i.e., leave the system as soon as they are created. Then using the relations

$$n_n = \frac{(\mu_n^2 - m_n^2)^{3/2}}{3\pi^2} \quad (5)$$

for neutron density. We have $n_B = n_p + n_n$, the total baryon number density.

Now in neutron stars the dynamic nature of chemical equilibrium among the constituents is mainly controlled by the following weak interaction processes (URCA and modified URCA processes [14]). These are also the most important cooling processes by neutrino / anti-neutrino emissions and are given by

$$\begin{aligned} n &\rightarrow p + e^- + \bar{\nu}_e \\ p + e^- &\rightarrow n + \nu_e \end{aligned} \quad (6)$$

Now from the solution of Dirac equation for electrons in presence of a quantizing magnetic field it can be shown that when only the zeroth Landau level is occupied by electrons, all the electrons will be aligned in the opposite direction to the magnetic field. Then we can write down the following relations from the condition of dynamical chemical equilibrium

$$\mu_{p\uparrow} = \mu_{p\downarrow}, \quad \mu_{n\uparrow} = \mu_{n\downarrow} \quad (7)$$

with

$$\begin{aligned} \mu_{n\uparrow} &= \mu_{p\uparrow} + \mu_{e\downarrow} \\ \mu_{n\downarrow} &= \mu_{p\downarrow} + \mu_{e\downarrow} = \mu_{p\uparrow} + \mu_{e\downarrow} \end{aligned}$$

Here we have indicated the direction of polarization of electrons by $\downarrow$. Which simply means that both the proton and neutron matter sectors are exactly spin symmetric. Although neutrinos / anti-neutrinos are assumed to be non-degenerate, they must carry the spin signature of the processes in which they are produced. This scenario of neutrino / anti-neutrino emissions are almost identical with the famous experiment by C.S. Wu using polarized Co^{60} . The only difference is that here instead of polarized nucleons, we have polarized electrons, which are either absorbed or emitted by the nucleons. To show, that the spin symmetric picture of both neutron and proton sector in a strongly magnetized neutron star is physically impossible we investigate the Pauli para-magnetism for neutron component. We are not considering the proton matter, since they may occupy some of their non-zero Landau levels and as a consequence the quantum dia-magnetism of proton matter may dominate over the Pauli para-magnetism. Now it is very easy to show that the total neutron number density in the relativistic region is given by

$$n_n = \frac{[2\mu_n^* B(2\mu_n^* B + 2m_n)]^{3/2}}{6\pi^2} (2x_n^{3/2} - 1) \quad (8)$$

where μ_n^* is the magnitude of neutron anomalous magnetic dipole moment and

$$x_n = \frac{\varepsilon_n^2(B) - m_n^2}{2\mu_n^* B(2\mu_n^* B + 2m_n)}, \quad (9)$$

$\varepsilon_n(B)$ is the single particle energy for neutron in presence of the magnetic field B , which is given by

$$\varepsilon_n(B) = \{m_n^2 + x_n[R_{3/2}]^{-2/3}(\varepsilon_0^2 - m_n^2)\}^{1/2} \quad (10)$$

where ε_0 is the value of $\varepsilon_n(B)$ in absence of magnetic field. Now we define

$$R_{3/2} = \frac{1}{2}[2x_n^{3/2} - 1] \quad (11)$$

then the density of up-spin neutrons is given by

$$n_n^\uparrow(B) = \frac{n_n}{2} x_n^{3/2} [R_{3/2}(x_n)]^{-1} \quad (12)$$

and similarly for the down-spin case we have

$$n_n^\downarrow(B) = n_n^\uparrow[1 - x_n^{-3/2}] \quad (13)$$

Now for $x = \infty$, which corresponds to field free case, then we have $n_n^\uparrow = n_n^\downarrow = n_n/2$, which indicates that both up and down spin neutron states are equi-probable. On the other hand for $x \rightarrow 1$, we have $n_n^\uparrow(B) = n_n$ and $n_n^\downarrow(B) = 0$, where this limiting condition indicates the complete saturation, the corresponding strength of magnetic field is given by

$$B_s = \frac{1}{2\mu_n^*} [\{(6\pi^2 n_n)^{2/3} + m_n^2\}^{1/2} - m_n] \quad (14)$$

and the single particle energy for neutron in the saturation limit is given by

$$\varepsilon^{(s)}(B = B_s) = [2^{2/3} \varepsilon_0^2 - (2^{2/3} - 1)m_n^2]^{1/2} \quad (15)$$

In fig.(2) we have plotted $n_n^\uparrow$ and $n_n^\downarrow$ for various values of baryon number densities having astrophysical relevance. We have obtained the corresponding magnetic field strength from fig.(1). In these studies, since we are interested only on those electrons occupying the zeroth Landau level, we have taken B from fig.(1) slightly above the $n_{\max}^{(e)} = 1$ line ($= 2.5 \times B$ of $n_{\max}^{(e)} = 1$ line). This figure shows that within the range of densities and magnetic fields of astrophysical relevance, neutron matter sector can not be spin symmetric ($n_n^\uparrow \neq n_n^\downarrow$).

We shall now go over to interacting picture of hadronic matter. We assume a $\sigma - \omega - \rho$ exchange type mean field interaction scenario. Then the chemical potentials are given by

$$\begin{aligned} \mu_p &= \epsilon_f^p + g_\omega \omega_0 + \frac{1}{2} g_\rho \rho_3^0 \\ \mu_n &= \epsilon_f^n + g_\omega \omega_0 - \frac{1}{2} g_\rho \rho_3^0 \\ \mu_e &= \epsilon_f^e \end{aligned} \quad (16)$$

where ω_0 and ρ_3^0 are the mean values for vector and the third component of iso-vector fields respectively, g_i 's are the corresponding coupling constants ($i = \sigma, \omega$ and ρ). Assuming the non-zero values for the magnetic moment for the constituents, the Fermi energies ϵ_f^i 's ($i = p, n$ and e) are then given by

$$\begin{aligned} \epsilon_f^p &= \left(p_f^2 + ((m^* + 2eB\nu)^{1/2} + \Delta_p s_p B)^2 \right)^{1/2} \\ \epsilon_f^n &= \left(p_f^2 + (m^* + \Delta_n s_n B)^2 \right)^{1/2} \\ \epsilon_f^e &= \left(p_f^2 + ((m^2 + 2eB\nu)^{1/2} + \Delta_e s_e B)^2 \right)^{1/2} \end{aligned} \quad (17)$$

where $m^* = m_n - g_\sigma \sigma_0$, the nucleon effective mass, Δ_i 's are related to the magnetic moments of the components, including the neutron part, $s_i = +1$ and -1 for spin up and spin down cases respectively. If we neglect the anomalous magnetic moments, we get back the results as presented for the non-interacting case, with bare nucleon mass replaced by the effective mass. In that case it is again very easy to show that the spin symmetric neutron-proton matter can not exist inside a strongly magnetized neutron star. On the other hand in presence of (anomalous) magnetic moments, either we must have

$$\mu_{p\uparrow} \neq \mu_{p\downarrow}, \quad \mu_{n\uparrow} \neq \mu_{n\downarrow} \quad (18)$$

instead of eqn.(7), or $\Delta_p = \Delta_n = 0$ to satisfy eqn.(7). It is obvious that this inequality is not consistent with the β -equilibrium condition. On the other hand the assumption of zero (anomalous) magnetic moments is against the

established quantum mechanical results. Hence in this case also the possibility of spin symmetric neutron matter or proton matter is questionable.

We therefore conclude that if the magnetic field strength inside a neutron star is such that the electrons occupy only the zeroth Landau level and are in β -equilibrium with neutron and proton components, then the necessary conditions of spin symmetry are not satisfied. Alternatively, we may conclude that the magnetic field strength can not go beyond 10^{17}G , so that electrons occupy only their zeroth Landau level.




FIG. 1: The curve separating only zero Landau level occupied region and non-zero Landau level occupied region for electrons.




FIG. 2: Variation of neutron number density $n_n^{(i)}$ (expressed in terms of n_n) for a spin either parallel or anti-parallel to the external magnetic field B , with the baryon number density n_B (expressed in fm^{-3}). The upper curve is for $n_n^{\uparrow}$ and lower one for $n_n^{\downarrow}$. In this case electrons are occupying only their zeroth Landau level.

-
- [1] R.C. Duncan and C. Thompson, *Astrophys. J. Lett.* **392**, L9 (1992); C. Thompson and R.C. Duncan, *Astrophys. J.* **408**, 194 (1993); C. Thompson and R.C. Duncan, *MNRAS* **275**, 255 (1995); C. Thompson and R.C. Duncan, *Astrophys. J.* **473**, 322 (1996).
 - [2] P.M. Woods et al., *Astrophys. J. Lett.* **519**, L139 (1999); C. Kouveliotou, et al., *Nature* **391**, 235 (1999).
 - [3] K. Hurley, et al., *Astrophys. Jour.* **442**, L111 (1999).
 - [4] S. Mereghetti and L. Stella, *Astrophys. Jour.* **442**, L17 (1999); J. van Paradijs, R.E. Taam and E.P.J. van den Heuvel, *Astron. Astrophys.* **299**, L41 (1995); S. Mereghetti, astro-ph/99111252; see also A. Reisenegger, astro-ph/01003010.
 - [5] D. Bandopadhyaya, S. Chakrabarty, P. Dey and S. Pal, *Phys. Rev.* **D58**, 121301 (1998); T. Ghosh and S. Chakrabarty, *Phys. Rev.* **D63**, 0403006 (2001); T. Ghosh and S. Chakrabarty, *Int. Jour. Mod. Phys.* **D10**, 89 (2001).
 - [6] S. Chakrabarty, D. Bandopadhyay and S. Pal, *Phys. Rev. Lett.* **78**, 2898 (1997); D. Bandopadhyay, S. Chakrabarty and S. Pal, *Phys. Rev. Lett.* **79**, 2176 (1997).
 - [7] C.Y. Cardall, M. Prakash and J.M. Lattimer, astro-ph/0011148 and references therein; E. Roulet, astro-ph/9711206; L.B. Leinson and A. Pérez, astro-ph/9711216; D.G. Yakovlev and A.D. Kaminkar, *The Equation of States in Astrophysics*, eds. G. Chabrier and E. Schatzman P.214, Cambridge Univ. Press, 1994.
 - [8] V.G. Bezchastrov and P. Haensel, astro-ph/9608090.
 - [9] Sutapa Ghosh, Soma Mandal and Somenath Chakrabarty, *Ann. Phys.* **312** (2004) 398.
 - [10] Soma Mandal, Sutapa Ghosh and Somenath Chakrabarty, *Phys. Rev.* **C74**, 015801 (2006).
 - [11] A. Melatos, *Astrophys. Jour.* **519**, L77 (1999); A. Melatos, *MNRAS* **313**, 217 (2000).
 - [12] S. Bonazzola et al, *Astron. & Astrophysics.* **278**, 421 (1993); M. Bocquet et al, *Astron. & Astrophysics.* **301**, 757 (1995).
 - [13] B. Bertotti and A.M. Anile, *Astron. & Astrophysics.* **28**, 429 (1973); C. Cutler and D.I. Jones, gr-qc/0008021; K. Konno, T. Obata and Y. Kojima, gr-qc/9910038; A.P. Martinez et al, hep-ph/0011399; M. Chaichian et al, *Phys. Rev. Lett.* **84**, 5261 (2000); Guangjun Mao, Akira Iwamoto and Zhuxia Li, astro-ph/0109221; A. Melatos, *Astrophys. Jour.* **519**, L77 (1999); A. Melatos, *MNRAS* **313**, 217 (2000); R. González Felipe et al, astro-ph/0207150 and references therein.
 - [14] S. Shapiro and S. Teukolsky, 1983, *Black Holes, White Dwarfs, and Neutron Stars*, John Wiley and Sons, Inc.